Homework Problem Set #4

Math 5467

March 20, 2006

Exercise 1 *Find the z-transform for the sequence* $\mathbf{x}(n) = 0$ *for* $n < 0$ *and* $\mathbf{x}(n) = (2/3)^n$ *for* $n \geq 0$, *i.e.,*

$$\mathbf{x} = \begin{pmatrix} \cdots & 0 & 0 & 1 & \frac{2}{3} & \frac{4}{9} & \cdots & \frac{2^n}{3^n} & \cdots \end{pmatrix}$$

Exercise 2 *Show explicitly that the double-shift orthogonality condition in the frequency domain*

$$\int_{-\pi}^{\pi} e^{2ik\omega} |C(\omega)|^2 d\omega = 2\pi\delta_{k0}$$

for integer k, *is equivalent to*

$$|C(\omega)|^2 + |C(\omega + \pi)|^2 = 2.$$

Exercise 3 *Plot the frequency response function* $|H(\omega)|$ *in the interval* $0 \leq \omega \leq 2\pi$ *for the Daubechies 4 tap filter. Point out the features of the graph that cause this filter to belong to the maxflat class.*

Exercise 4 *Let* $\phi(t)$ *and* $w(t)$ *be the Haar scaling and wavelet functions. Let* V_j *and* W_j *be the spaces generated by* $\phi_{j,k}(t) = 2^{j/2}\phi(2^j t - k)$ *and* $w_{j,k}(t) = 2^{j/2} w(2^j t - k)$, $k = 0, \pm 1, \cdots$, *respectively. Let* $f(t)$ *be defined on* $0 \leq t < 1$ *and given by*

$$f(t) = \begin{cases} 4 & 0 \leq t < 1/4 \\ 1 & 1/4 \leq t < 1/2 \\ -1 & 1/2 \leq t < 3/4 \\ 2 & 3/4 \leq t < 1. \end{cases}$$

1. *Express f in terms of the basis for V_2.*

2. *Decompose f into its component parts in W_1, W_0, and V_0. In other words, find the Haar wavelet decomposition for f.*

3. *Sketch each of the four decompositions.*

Exercise 5 *Let $\phi(t)$ and w be the Haar scaling and wavelet functions, respectively. Let V_j and W_j be the spaces generated by $\phi_{j,k}(t) = 2^{j/2}\phi(2^j t - k)$ and $w_{j,k}(t) = 2^{j/2}w(2^j t - k)$, $k = 0, \pm 1, \cdots$, respectively. Suppose the real valued function $f(t) = \sum_k a_k \phi_{1,k}(t)$ belongs to V_1 and $f \perp V_0$. Show explicitly that $a_{2\ell+1} = -a_{2\ell}$ for all integers ℓ. Conclude that f can be expressed as a linear combination of the functions, $w_{0,k}(t)$, hence that $f \in W_0$.*