

**GENESIS OF GROUPS AND ITS RAMIFICATION FROM
n-LATERAL MULTIPLE HYPERGEOMETRIC FUNCTION**

By


Anand Singh

and

H.S. Dhami

IMA Preprint Series # 1605

February 1999



INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS

UNIVERSITY OF MINNESOTA

514 Vincent Hall

206 Church Street S.E.

Minneapolis, Minnesota 55455-0436

Phone: 612/624-6066 Fax: 612/626-7370

URL: <http://www.ima.umn.edu>

GENESIS OF GROUPS AND ITS RAMIFICATION FROM n- LATERAL MULTIPLE HYPERGEOMETRIC FUNCTION

Anand Singh and H.S. Dhami*

Department of Mathematics

University of Kumaun

Almora Campus,

Almora (U.P) 263601 India

In an attempt to narrow down the gap between theory and applications, the two advanced branches of mathematics, special function and group theory have been brought closer by formation of finite dimensional group for n- lateral multiple hypergeometric functions in our earlier study. Here efforts are being made to generate subgroups, ring homomorphism and isomorphism have been also established for B- function involving two variables.

Key words :- Limit Point / Lauricella function / Homomorphism / identity element.

1. INTRODUCTION

The works of Vilinkin³ and Talman² had provided with a basement for derivation of different special functions of one variable and their properties from the theory of group representations. We, in our earlier work¹ have constructed finite multiplicative group for n- lateral multiple hypergeometric functions.

Here an attempt is being made to go ahead in the direction of establishing further relationship between special functions and group theory by generation of subgroups, cosets, normal subgroup, homomorphism, isomorphism, ring and topological space for our generalized B- function involving two variables.

* To whom all correspondence be mailed

2. GENERAL FORMULATION

From the group (S_.) generated by result (3.7) of our earlier study, we can select subgroups of following nature,

- (a). Where parameters of later half are zero.

As an example we may cite,

$$S_1 = \{B^{10}, B^{20}, B^{30}\} \quad \text{.....(2.1)}$$

Whose first and third elements are inverse of each other and second element is identity .

- (b). Where parameters of first half are zero.

As an example, the following subgroup may be mentioned

$$S_2 = \{B^{01}, B^{02}, B^{03}\} \quad \text{.....(2.2)}$$

Here again the first and third elements are inverse of each other and second element is identity.

- (c). Where parameters of both half are considered, the subgroup of order 3 in this case shall be

$$S_3 = \{B^{11}, B^{22}, B^{33}\} \quad \text{.....(2.3)}$$

While the subgroup of order 5 shall be

$$S_4 = \{B^{12}, B^{21}, B^{23}, B^{32}, B^{22}\} \quad \text{.....(2.4)}$$

the last element of this subgroup shall be the identity element while first & fourth and second & third elements (taken in pairs) shall be inverse of each other.

Here it is evident that subgroups formed in present case do not follow the general rule (identities of group and its subgroups are same), due to the reason that parameters of one of the two halves have been omitted. So in this case the identity of the group shall be equal to the product of the identities of subgroups.

3.COMPOSITION OF COSETS

Considering the subgroup S_3 (given by equation (2.3)) we can form left and right cosets as,

$$\begin{aligned} S_3 * B^{01} &= \{ B^{11} * B^{01}, B^{22} * B^{01}, B^{33} * B^{01} \} \\ &= \{ B^{11}, B^{22}, B^{32} \} \end{aligned} \dots\dots\dots(3.1)$$

and

$$\begin{aligned} B^{01} * S_3 &= \{ B^{01} * B^{11}, B^{01} * B^{22}, B^{01} * B^{33} \} \\ &= \{ B^{11}, B^{22}, B^{32} \} \end{aligned} \dots\dots\dots(3.2)$$

Which implies the condition required for our subgroup to be a normal subgroup.

4. ISOMORPHISM OF SUBGROUPS

Let $f : S \rightarrow S$ be defined as

$$f(B^{pq}) = B^{qp} \dots\dots\dots(4.1)$$

then it will be proved that f is an isomorphism.

The mapping is one -one onto from mapping defined by relationship (4.1) and homomorphism can be depicted from the following exemplary case,

$$\begin{aligned} f(B^{10} * B^{03}) &= f(B^{13}) = B^{31} \\ &= B^{01} * B^{30} = f(B^{10}) * f(B^{03}) \end{aligned}$$

5. FORMATION OF RING

Let us incorporate the element B^{00} (normal B-function) in the Set S of multiplicative group $(S, *)$ obtained in section 3 (result (3.7)) of our earlier work and to binary operators \oplus and \odot be defined as

$$\begin{aligned} B^{ij} \oplus B^{pq} &= \text{products of non repeated terms of } B^{ij} \text{ and } B^{pq} \\ B^{ij} \odot B^{pq} &= \text{products of repeated terms of } B^{ij} \text{ and } B^{pq} \end{aligned}$$

then (S, \oplus) shall be an abelian group and (S, \odot) will be a semi-group.

Distributive Law shall hold good in this case as evident from the following example

$$B^{12} \odot [B^{23} \oplus B^{13}] = (B^{12} \odot B^{23}) \oplus (B^{12} \odot B^{13})$$

Here

$$B^{12} \odot [B^{23} \oplus B^{13}] = B^{12} \odot B^{30} = B^{00} \dots\dots\dots(5.2)$$

and

$$(B^{12} \odot B^{23}) \oplus (B^{12} \odot B^{13}) = B^{13} \oplus B^{13} = B^{00} \dots\dots\dots(5.3)$$

Which demonstrates that ring shall be a ring with zero divisor. B^{00} shall be the zero of the ring and B^{22} shall be its unity element.

6. TOPOLOGICAL SPACE

Let us defined the set

$$X = \{ B^{10}, B^{20}, B^{30} \} \dots\dots\dots(6.1)$$

then the different topologies for X can be formed.

The smallest topology for X shall be

$$\mathfrak{T}_1 = \{ B^{00}, X \} \dots\dots\dots(6.2)$$

B^{00} shall be taken as null set.

The largest topology on X shall be

$$\mathfrak{T}_{14} = \{ B^{00}, \{B^{10}\}, \{B^{20}\}, \{B^{30}\}, \{B^{10}, B^{20}\}, \{B^{20}, B^{30}\}, \{B^{10}, B^{30}\}, X \} \dots\dots\dots(6.3)$$

Neighbourhood system for every element can be seen as it is evident from the following example

$N(B^{10})$ in \mathfrak{T}_{14} shall be

$$\{ \{B^{10}\}, \{B^{10}, B^{20}\}, \{B^{10}, B^{30}\}, X \}$$

In the smallest topological space (X, \mathfrak{T}_1)

If we consider set $A = \{B^{10}, B^{20}\}$

then each point of X shall be a limit point for A and the derived set in this case shall be X itself.

REFERENCES

1. Anand Singh & H.S.Dhami (1998) Formation of finite multiplicative group for n-lateral multiple hypergeometric functions, IMA Preprint series #1596.
2. James, Talman (1968) Special functions : A group theoretic approach, W.A.Benjamin, Inc. New York.
3. N.Y. Vilinkin (1968) Special functions and the theory of group representations AMS Transl., Providence, Rhode Island. [English Transl.(1968)].