

**ON THE EXISTENCE OF CERTAIN SMOOTH TORIC VARIETIES**

By

**Jörg Gretenkort  
Peter Kleinschmidt  
and  
Bernd Sturmfels**

**IMA Preprint Series # 354**

October 1987

**INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS**

**UNIVERSITY OF MINNESOTA**

514 Vincent Hall  
206 Church Street S.E.  
Minneapolis, Minnesota 55455

## Recent IMA Preprints (Continued)

#	Author(s)	Title	#	Author(s)	Title
241	Harry Kesten	Scaling Relations for 2D-Percolation	242	A. Leizarowitz	Infinite Horizon Optimization for Markov Processes with Finite States Spaces
243	Louis H. Y. Chen	The Rate of Convergence in a Central Limit Theorem for Dependent Random Variables with Arbitrary Index Set	244	G. Kallianpur	Stochastic Differential Equations in Duals of Nuclear Spaces with some Applications
245	Tzuu-Shuh Chiang, Yunshong Chow, Yuh-Jia Lee	Evaluation of Certain Functional Integrals	246	L. Karp, M. Pinsky	The First Eigenvalue of a Small Geodesic Ball in a Riemannian Manifold
247	Chi-Sing Man	Towards an Acoustoelastic Theory for Measurement of Residual Stress	248	Andreas Stoll	Invariance Principles for Brownian Intersection Local Time and Polymer Measures
249	R. W. R. Darling	Rate of Growth of the Coalescent Set in a Coalescing Stochastic Flow	250	R. Cohen, R. Hardt, D. Kinderlehrer, S. Y. Lin, M. Luskin	Minimum Energy for Liquid Crystals: Computational Results
251	Suzanne M. Lenhart	Viscosity Solutions for Weakly Coupled Systems of First Order PDEs	252	M. Cranston, E. Fabes, Z. Zhao	Condition Gauge and Potential Theory for the Schrödinger Operator
253	H. Brezis, J. M. Coron, E. H. Lieb	Harmonic Maps with Defects	254	A. Carverhill	Flows of Stochastic Dynamical Systems: Nontriviality of the Lyapunov Spectrum
255	A. Carverhill	Conditioning a 'Lifted' Stochastic System in a Product Case	256	R. J. Williams	Local Time and Excursions of Reflected Brownian Motion
257	H. Folmer, S. Orey	Large Deviations for the Empirical Field of a Gibbs Measure	258	A. Leizarowitz	Characterization of Optimal Trajectories on an Infinite Horizon
259	Y. Giga, T. Miyakawa, H. Osada	Two Dimensional Navier Stokes Flow with Measures as Initial Vorticity	260	M. Chipot, V. Oliker	Sur une propriété Des Fonctions Propres De l'opérateur De Laplace Beltrami
261	V. Perez-Abreu	Decompositions of Semimartingales on Duals of Countably Nuclear Spaces	262	J. M. Ball	Does Rank-One Convexity Imply Quasiconvexity?
263	B. Cockburn	The Quasi-Monotone Schemes for Scalar Conservation Laws. Part 1	264	K. A. Pericak-Spector	On Radially Symmetric Simple Waves in Elasticity
265	P. N. Shivakumar, Chi-Sing Man, Simon W. Rabkin	Modelling of the Heart and Pericardium at End-Diastole	266	Jose-Luis Menaldi	Probabilistic View of Estimates for Finite Difference Methods
267	Robert Hardt, Harold Rosenberg	Open Book Structures and Unicity of Minimal Submanifolds	268	Bernardo Cockburn	The Quasi-Monotone Schemes for Scalar Conservation Laws
269	H. R. Jauslin, W. Zimmermann, Jr.	Dynamics of a Model for an Ac Josephson Effect in Superfluid 4He	270	A. K. Kapila	Introductory Lecture on Reacting Flows
271	J. C. Taylor	Do Minimal Solutions of Heat Equations Characterize Diffusions?	272	J. C. Taylor	The Minimal Eigenfunctions Characterize the Ornstein-Uhlenbeck Process
273	Chi-Sing Man, Quan-Xin Sun	On the Significance of Normal Stress Effects in the Flow of Glaciers	274	Omar Hijab	On Partially Observed Control of Markov Processes
275	Lawrence Gray	The Behavior of Processes with Statistical Mechanical Properties	276	R. Hardt, D. Kinderlehrer, M. Luskin	Remarks about the Mathematical Theory
277	Cockburn, B.	The Quasi-Monotone Schemes for Scalar Conservation Laws Part III	278	M. Chipot, T. Sideris	On the Abelian Higgs Model
279	C. Foias, B. Nicolaenko, G.R.Sell, R. Temam	Inertial Manifolds for the Kuramoto-Sivashinsky Equation and an Estimate of their Lowest Dimension	280	R. Duran	On the Approximation of Miscible Displacement in Porous Media by a Method of Characteristics Combined with a Mixed Method
281	H. Aixiang, Zhang Bo	The Convergence for Nodal Expansion Method	282	V. Tverzky	Dispersive Bulk Parameters for Coherent Propagation in Correlated Random Distributions
283	W.Th.F. den Hollander	Mixing Properties for Random Walk in Random Scenery	284	H.R. Jauslin	Nondifferentiable Potentials for Nonequilibrium Steady States
285	K. Meyer, G.R.Sell	Homoclinic Orbits and Bernoulli Bundles in Almost Periodic Systems	286	J. Douglas, Jr., Y. Yuan	Finite Difference Methods for the Transient Behavior of a Semiconductor Device
287	Li Kaitai, Yan Ningning	The Extrapolation for Boundary Finite Elements	288	R. Durrett, R.H. Schonmann	Stochastic Growth Models
289	David Kinderlehrer	Remarks about Equilibrium Configurations of Crystals	290	D.G. Aronson, J.L. Vazquez	Eventual $C^\infty$ -Regularity and Concavity for Flows in One-Dimensional Porous Media
291	L.R. Scott, J.M. Boyle, B. Bagheri	Distributed Data Structures for Scientific Computation	292	J. Douglas, Jr., P.J. Paez Leme, T. Arbogast, T. Schmitt	Simulation of Flow in Naturally Fractured Petroleum Reservoirs
293	D.G. Aronson, L.A. Caffarelli	Optimal Regularity for One-Dimensional Porous Medium Flow	294	Haim Brezis	Liquid Crystals and Energy Estimates for $S^2$ -Valued Maps
295	Todd Arbogast	Analysis of the Simulation of Single Phase Flow Through a Naturally Fractured Reservoir	296	He Yinnian, Li Kaitai	The Coupling Method of Finite Elements and Boundary Elements for Radiation Problems
297	T. Cazenave, A. Haraux, L. Vazquez, F.B. Weissler	Nonlinear Effects in the Wave Equation with a Cubic Restoring Force	298	M. Chipot, F.B. Weissler	Some Blow Up Results for a Nonlinear Parabolic Equation with a Gradient Term
299	Li Kaitai	Perturbation Solutions of Simple and Double Parabolic Equation with a Gradient Term	300	Chen Zhanqin, Li Kaitai	The Convergence on the Multigrid Algorithm for Navier-Stokes Equations
301	A. Gerardi, G. Nappo	Martingale Approach for Modeling DNA Synthesis	302	D.N. Arnold, L. Ridgway Scott, Michael Vogelius	Regular Inversion of the Divergence Operator with Dirichlet Boundary Conditions on a Polygon
303	Duran, Ricardo G.	Error Analysis in $L^p$ , $1 \leq p \leq \infty$ , for Mixed Definite Element Methods for Linear and Quasi-Linear Elliptic Problems	304	Nochetto, Ricardo H., Verdi, Claudio	An Efficient Linear Scheme to Approximate Parabolic Free Boundary Problems: Error Estimation and Implementation

# On the existence of certain smooth toric varieties

JÖRG GRETENKORT, PETER KLEINSCHMIDT<sup>†</sup>, AND BERND STURMFELS

**Abstract.** We prove that the combinatorial types of those cone systems which correspond to complete smooth toric varieties are more restricted than for complete toric varieties: The toric varieties corresponding to essentially all types of cyclic polytopes possess singularities. This yields a negative answer to a problem stated by G. Ewald. Some consequences and problems concerning Mathematical Programming and the rational cohomology of smooth toric varieties are discussed.

## 1. Introduction and basic terminology.

Recently, there has been much interaction between the combinatorial theory of polytopes and the theory of toric varieties [1,3,7,8,10]. A *toric  $d$ -variety* is a  $T$ -invariant subvariety of a  $T$ -invariant completion of the torus  $T = (k^*)^d$  ( $k$  can be  $\mathbb{C}$  or any algebraically closed field). It can be conveniently described in terms of a *fan*, i.e. a finite system of cones in  $\mathbb{R}^d$  spanned by integer lattice points. In this paper, we will not explain the numerous correspondences between properties of fans and properties of toric varieties. The interested reader should consult [2] and [8] for details.

Here, we will concentrate on those fans that correspond to complete smooth toric  $d$ -varieties.

Let  $\Delta$  be a simplicial  $(d-1)$ -complex with the following properties:

- (1)  $\Delta$  is embedded in  $\mathbb{R}^d$ ,
- (2) the underlying set of  $\Delta$ , denoted  $|\Delta|$ , is homeomorphic to the unit-sphere  $S^{d-1}$ ,
- (3) the bounded component of  $\mathbb{R}^d$  whose boundary is  $|\Delta|$  contains the origin in its interior and is star-shaped with respect to the origin.
- (4) the 0-cells (vertices) of  $\Delta$  are primitive lattice points in  $\mathbb{Z}^d$ ,
- (5) let  $x_1, \dots, x_d$  be the vertices of a  $(d-1)$ -cell of  $\Delta$ , then  $|\det(x_1, \dots, x_d)| = 1$ .

Let  $\Sigma_\Delta$  be the system of those cones whose apex is the origin and whose bases are the cells of  $\Delta$ . We call  $\Sigma_\Delta$  a *complete regular fan*. In the absence of property (5) we call it a *complete fan*.

It is well known that all complete smooth toric  $d$ -varieties can be described in terms of complete regular fans [2] (smoothness is equivalent to condition (5)). Let  $X_\Sigma$  denote the variety corresponding to the fan  $\Sigma$ . Two fans  $\Sigma$  and  $\Sigma'$  are *combinatorially equivalent* (or isomorphic) if the complexes  $\Delta$  and  $\Delta'$  which define  $\Sigma$  and  $\Sigma'$  are combinatorially equivalent, i.e. there is a bijective inclusion preserving map from  $\Delta$  to  $\Delta'$ .

---

<sup>†</sup>Supported in part by the Institute for Mathematics and its Applications, University of Minnesota, Minneapolis, Minnesota

The varieties  $X_\Sigma$  and  $X_{\Sigma'}$  are isomorphic if and only if such a combinatorial equivalence for  $\Sigma$  and  $\Sigma'$  can be induced by a unimodular transformation of  $\mathbb{R}^d$ . So, the study of toric varieties can be viewed as the study of invariants of fans under unimodular transformations.

At the 4th Geometry Symposium in Siegen, G. Ewald [4] posed the following question: *Let  $\Delta$  be a simplicial  $(d - 1)$ -dimensional spherical complex. Then, does there exist a spherical complex  $\Delta'$  combinatorially equivalent to  $\Delta$  such that  $\Delta'$  determines a complete regular fan?*

This question was answered positively in [7] for the case that the number of vertices of  $\Delta$  does not exceed  $d + 2$ . In the present paper we give an infinite family of counterexamples to this question. The smallest complex in our family has  $d + 3$  vertices and is therefore minimal with the desired property.

**THEOREM.** *Let  $\Delta$  be a complex which is isomorphic to the boundary-complex of the cyclic 4-polytope with  $n \geq 7$  vertices (see [5] for the definition of cyclic polytopes). Then there does not exist a complex  $\Delta'$  with the properties (1)-(5) which is combinatorially equivalent to  $\Delta$ .*

This Theorem implies that no complete smooth toric 4-variety has an underlying cone-system stemming from a cyclic 4-polytope with more than 6 vertices. Hence, the existence of certain smooth toric varieties may be ruled out on grounds of the combinatorial type of a fan alone. In contrast to this it is well known [10] that every combinatorial type of a simplicial polytope gives rise to (many) complete fans which correspond to projective toric varieties. However, our result shows that under the regularity assumption this may not hold.

The proof of the Theorem proceeds in two steps. In Section 2 we will establish the case  $n = 7$ , and in Section 3 a different technique of proof is used to settle the case  $n \geq 8$ . Finally, in Section 4 we will discuss some consequences and problems related to our result.

## 2. On the cyclic 4-polytope with 7 vertices.

Let us assume that  $\Delta$  is a complex which has properties (1) - (5) of the last section and which is isomorphic to the boundary complex of the cyclic 4-polytope with 7 vertices. Let  $1, 2, \dots, 7$  denote the vertices of  $\Delta$ . Then the following table lists the 14 tetrahedra which are the maximal cells of  $\Delta$ .

A = 1245	E = 1347	I = 2356	M = 2467
B = 1246	F = 1356	J = 2357	N = 3467
C = 1256	G = 1357	K = 2367	
D = 1346	H = 1457	L = 2457	

Table 1

Let  $i_1, \dots, i_4$  be the vertices of one of the listed tetrahedra and let  $i_1, i_2, i_3, i_5$  be the vertices of the unique tetrahedron in the list with which the former shares the vertices  $i_1, i_2$  and  $i_3$ . Then it follows from (5) that the following linear relation holds:

$$(*) \quad i_4 + i_5 + \sum_{j=1}^3 \lambda_j i_j = 0 \quad (\text{for some integers } \lambda_j).$$

As  $\Delta$  has 28 triangles, each of which is contained in precisely 2 of the tetrahedra, there are exactly 28 linear relations of type (\*). (In [7] and [8] extensive use of these relations has been made for the classification of toric varieties).

Interpreting the coefficients of the points  $i_1, \dots, i_7$  in a relation of type (\*) as vectors in  $\mathbf{R}^7$  (we call those vectors *linear dependencies*), it follows that the linear span of the linear dependencies is only 3-dimensional.

So, there is a lot of redundancy in the 28 relations of type (\*). Using just this fact, it is easy to check that every relation (\*) is equal to one of the 14 relations listed by their coefficients in Table 2. The symbols in the left column are the tetrahedra from which the coefficients of the relation in the respective row come. The numbers  $\lambda_1, \lambda_2, \dots, \lambda_{35}$  are undetermined coefficients.

Vertices Tetrahedra	1	2	3	4	5	6	7
AB	$\lambda_1$	$\lambda_2$	0	1	1	1	0
AL	1	1	0	$\lambda_3$	$\lambda_4$	0	1
BD	$\lambda_5$	1	1	$\lambda_6$	0	$\lambda_7$	0
BM	1	$\lambda_8$	0	$\lambda_9$	0	$\lambda_{10}$	1
CF	1	1	1	0	$\lambda_{11}$	$\lambda_{12}$	0
DE	1	0	$\lambda_{13}$	$\lambda_{14}$	0	1	1
DF	$\lambda_{15}$	0	$\lambda_{16}$	1	1	$\lambda_{17}$	0
FG	$\lambda_{18}$	0	$\lambda_{19}$	0	$\lambda_{20}$	1	1
GH	$\lambda_{21}$	0	1	1	1	0	$\lambda_{22}$
GJ	1	1	$\lambda_{23}$	0	$\lambda_{24}$	0	$\lambda_{25}$
JK	0	$\lambda_{26}$	$\lambda_{27}$	0	1	1	1
JL	0	$\lambda_{28}$	1	1	$\lambda_{29}$	0	$\lambda_{30}$
KN	0	1	1	1	0	$\lambda_{31}$	$\lambda_{32}$
LM	0	$\lambda_{33}$	0	$\lambda_{34}$	1	1	$\lambda_{35}$

Table 2

Using the fact that the rows  $AB$ ,  $BD$  and  $BM$  in the above table are linearly independent and hence span the space of all linear relations, we can deduce further relations between the various  $\lambda_i$ . This could be done systematically using affine Grassmann-Plücker relations. However, as we are satisfied when a contradictory relation is reached, we will not present all relations and details here. These can easily be checked.

From the representation of all relations as linear combinations of  $AB$ ,  $BD$  and  $BM$  we obtain among many other relations the following:

$$\begin{aligned} (1) \quad & \lambda_1 \cdot \lambda_{10} = 0 \\ (2) \quad & \lambda_2 \cdot \lambda_6 = 0 \\ (3) \quad & \lambda_5 \cdot \lambda_8 = 0 \end{aligned}$$

As in each relation at least one of the  $\lambda_i$  is equal to zero, we may branch the determination of all  $\lambda_i$  into 8 cases where 3 of the  $\lambda_i$  from (1), (2) and (3) are set to zero. Each of these cases yields a contradiction for some of the  $\lambda_i$ . For example, if  $\lambda_1 = \lambda_2 = \lambda_5 = 0$ , it follows (from the additional relation  $\lambda_5 - \lambda_1\lambda_6 = 1$ ) that  $1 = 0$ . This proves the case  $n = 7$  of the Theorem.

### 3. On cyclic 4-polytopes with 8 or more vertices.

In this section we complete the proof of the Theorem by showing that no cyclic 4-polytope with eight or more vertices gives rise to a smooth toric variety. Let  $\Delta$  be the boundary complex of a cyclic 4-polytope with  $n \geq 8$  vertices, and assume that  $\Delta$  is embedded with vertices  $x_1, x_2, \dots, x_n$  in  $\mathbb{R}^4$  such that the properties (1) – (5) in Section 1 are satisfied.

Every facet  $\{i, j, k, l\}$  of  $\Delta$  corresponds to a basis  $\{x_i, x_j, x_k, x_l\}$  of  $\mathbb{R}^4$ , and the orientations of all such ordered bases induce an orientation of the simplicial complex  $\Delta$ . The 3-sphere  $\Delta$  being closed, connected and orientable, there exists (up to inversion) a unique such intrinsic orientation  $\text{sign}_\Delta$ . This implies the following stronger version of property (5).

(5') Let  $x_i, x_j, x_k, x_l$  be the vertices of a tetrahedron of  $\Delta$ , then  $[i, j, k, l] = \text{sign}_\Delta(i, j, k, l)$ . Here and throughout this section we use the abbreviation  $[i, j, k, l] := \det(x_i, x_j, x_k, x_l)$ .

Consider the simplicial 3-complex  $\tilde{\Delta}$  on  $\{1, 2, \dots, 8\}$  which is defined by the following set of 15 tetrahedra:

$$\{ \{r, r+1, r+s, r+s+1\} \mid 1 \leq r \leq 5, 2 \leq s \leq 7-r \}.$$

By Gale's evenness condition [5],  $\tilde{\Delta}$  is a subcomplex of  $\Delta$ , and it is easy to check that  $\text{sign}_\Delta(r, r+1, r+s, r+s+1) = +1$  for all  $1 \leq r \leq 5$  and  $2 \leq s \leq 7-r$ . Observe that  $\tilde{\Delta}$  is a 3-ball, and therefore the unique orientation  $\text{sign}_{\tilde{\Delta}}$  of  $\tilde{\Delta}$  is induced from  $\text{sign}_\Delta$ .

Hence condition (5') implies

(5'')  $[r, r+1, r+s, r+s+1] = +1$  for all  $1 \leq r \leq 5$  and  $2 \leq s \leq 7-r$ .

We will show that there do not exist vectors  $x_1, x_2, \dots, x_8 \in \mathbb{R}^4$  satisfying the 15 determinant equations given in (5''). Note that the corresponding system of equations has a solution for  $n = 7$ , and hence the proof of the Theorem had to split into two parts.

Following the general philosophy of ‘‘Computational Synthetic Geometry’’ as outlined in [11], we shall give a compactly encoded nonrealizability proof for the synthetic geometry problem (5’). The set  $\{2, 3, 6, 7\}$  being a tetrahedron of  $\tilde{\Delta}$ , it can be assumed that  $(x_2, x_3, x_6, x_7)$  is a (positively oriented) orthonormal basis of  $\mathbf{R}^4$ , and we can write

$$(x_1, x_2, \dots, x_8) = \begin{pmatrix} a & 1 & 0 & e & i & 0 & 0 & m \\ b & 0 & 1 & f & j & 0 & 0 & n \\ c & 0 & 0 & g & k & 1 & 0 & o \\ d & 0 & 0 & h & l & 0 & 1 & p \end{pmatrix},$$

where  $a, b, c, \dots, o, p$  are indeterminants. Let

$$\begin{aligned} P_1 &:= 1 - [1234], & P_2 &:= 1 - [1245], & P_3 &:= 1 - [1256], & P_4 &:= 1 - [1267], & P_5 &:= 1 - [1278], \\ P_6 &:= 1 - [2345], & P_7 &:= 1 - [2356], & P_8 &:= 1 - [2367], & P_9 &:= 1 - [2378], & P_{10} &:= 1 - [3456], \\ P_{11} &:= 1 - [3467], & P_{12} &:= 1 - [3478], & P_{13} &:= 1 - [4567], & P_{14} &:= 1 - [4578], & P_{15} &:= 1 - [5678]. \end{aligned}$$

For example, we have  $P_{10} = 1 - el + hi$ .

Let  $I$  denote the ideal in  $\mathbf{Q}[a, b, \dots, p]$  generated by the polynomials  $P_1, P_2, \dots, P_{15}$ . It is clearly sufficient to show that  $I$  has no complex zeros. To prove this constructively, we shall give a *final polynomial* [11, Chapter 4] for the realizability problem in question, that is, we shall construct polynomials  $Q_1, \dots, Q_{15} \in \mathbf{Q}[a, b, \dots, p]$  such that  $\sum_{i=1}^{15} Q_i P_i = 1$ .

Consider the elements  $R_1$  and  $R_2$  of  $I$  which are defined as  $R_1 := eP_7 + P_{10} - P_{11}$ , and

$$\begin{aligned} R_2 &:= \\ &-n(c + k + icf)P_1 + inP_2 + (cn + kn + ign)P_3 + (o + inhk + ncl + nkl)P_4 \\ &+ P_5 - (m + 2in - ncd - nkd - ncdif + ifm + mfk)P_6 \\ &-(nk + nc + mg + 2ing - nicf - ndg(c + k + icf) + fmg(i + k) - cfe - enk)P_7 \\ &+(e - 1 + f(i + ie + ke))P_9 + (cf + nk)P_{10} - (1 + j - jm + f(k + c + i) + njd(c + k))P_{11} \\ &+(1 + fi + fk + j)P_{12} + (m - 1 - ncd - nkd)P_{13} + P_{14} - P_{15}. \end{aligned}$$

It can be seen that  $R_1 = ih$ , and that  $R_2 + 1$  is divisible by  $h$ . Let  $S_2 := (R_2 + 1)/h \in \mathbf{Q}[a, b, \dots, p]$ .

The combinatorial symmetry  $\sigma = (18)(27)(36)(45)$  of  $\tilde{\Delta}$  induces the symmetry  $\tau = (ap)(bo)(cn)(dm)(el)(fk)(gj)(hi)$  on the set  $\{P_1, \dots, P_{15}\}$  of generators of  $I$ . Applying the symmetry  $\tau$  to the above representation of  $R_2$  we get a representation of  $R_3 := \tau(R_2)$  as linear combination of the  $P_i$ . Clearly,  $R_3 + 1$  is divisible by  $i = \tau(h)$ , and we can set  $S_3 := (R_3 + 1)/i \in \mathbf{Q}[a, b, \dots, p]$ . Finally, observe the identity

$$S_2 S_3 R_1 - R_2 - h S_2 R_3 = 1.$$

This proves that  $1 \in I$ , and, moreover, with the above representations of  $R_1, R_2$  and  $R_3$ , we immediately obtain polynomials  $Q_i$  satisfying the desired identity  $\sum_{i=1}^{15} Q_i P_i = 1$ .

#### 4. Consequences and problems.

Let  $X_\Sigma$  be a complete toric  $d$ -variety given by a fan  $\Sigma$  or, equivalently, by a spherical complex  $\Delta$  with  $n$  vertices. Let  $f_i$  be the number of  $i$ -dimensional cells of  $\Delta$  and, for  $0 \leq i \leq d$ , let  $h_i$  denote the following quantities.

$$h_i := \sum_{j=0}^i \binom{d-j}{d-i} (-1)^{i-j} f_{j-1} \quad (f_{-1} := 1).$$

It is well known [9] that these  $h_i$  count the cocycles which generate the rational cohomology of  $X_\Sigma$ . More precisely,  $h_i = rk H^{2i}(X_\Sigma, \mathbb{Q})$ ,  $0 \leq i \leq d$  (note that the odd dimensional cohomology vanishes).

As  $|\Delta|$  is a sphere, the ‘‘Upper-bound-theorem for spheres’’ (‘‘UBTS’’ for short) [9] says that  $h_i \leq \binom{n-d-1}{i}$ , or (as  $h_1 = n - d$ )

$$h_i \leq \binom{h_1 + i - 1}{i}, \quad \text{for } 0 \leq i \leq [d/2].$$

The Dehn-Sommerville equations [5] are equivalent to  $h_i = h_{d-i}$ ,  $0 \leq i \leq d$ , so that the above bounds for the  $h_i$  imply bounds for the other  $h_i$  as well. All these bounds are simultaneously achieved if  $\Delta$  is isomorphic to the boundary complex of the cyclic  $d$ -polytope with  $n$  vertices.

As mentioned before, it is well known that for every combinatorial type of a simplicial  $d$ -polytope there is a complete (even projective) toric  $d$ -variety whose fan is given by a complex which is isomorphic to the boundary complex of that polytope. So, the above bounds for the  $h_i$  are also sharp upper bounds for the ranks of rational cohomology of complete toric  $d$ -varieties. So, UBTS has the following counterpart for toric varieties (call it ‘‘UBTV’’ for short):

Let  $X$  be a complete toric  $d$ -variety with  $rk H^2(X, \mathbb{Q}) =: k$  fixed. Then the following holds:

$$rk H^{2i}(X, \mathbb{Q}) \leq \binom{k + i - 1}{i}, \quad 0 \leq i \leq [d/2]$$

and the bounds are achieved by fans which are given by the boundary complexes of cyclic  $d$ -polytopes with  $k + d$  vertices. So the number  $k$  already determines sharp upper bounds for the number of higher dimensional cohomology classes.

It is a consequence of our Theorem in Section 1 that the statement about sharpness of the UBTV is not valid for *smooth*  $X$  with  $d = 4$  and  $k = 3$ . For the combinatorial type of a corresponding sphere with  $h_2 = 6$  (for sharpness) would have to be the one of the cyclic 4-polytope with 7 vertices [5]. Our Theorem, however implies that this type does not give rise to a smooth toric variety. So we have (in contrast to the UBTV):

COROLLARY 1. *Let  $X$  be a complete smooth toric 4-variety with  $\text{rk } H^2(X, \mathbb{Q}) = 3$ . Then  $\text{rk } H^4(X, \mathbb{Q}) \leq 5$ . This bound can be achieved.*

This corollary is an immediate consequence of our Theorem and the UBTS. The statement about sharpness follows by blowing-up a suitable 4-variety from [7]. It is not surprising that smoothness implies fewer cocycles. However, we are not aware of any other result which directly implies Corollary 1.

It would be of interest to prove a correct version of the UBTV for the smooth case, i.e., to determine sharp upper bounds for  $\text{rk } H^{2i}(X, \mathbb{Q})$  given  $\text{rk } H^2(X, \mathbb{Q})$  for complete smooth toric  $d$ -varieties. It may be guessed that asymptotically these bounds are by far smaller than those given in the UBTV. A result of this type would allow to rule out smoothness of a variety by computing some of the  $h_i$ .

Such bounds would be of interest in Mathematical Programming: One can interpret the cones of a complete fan as the cones spanned by the normals of the hyperplanes which determine a basic feasible solution of a nondegenerate LP [6]. The regularity condition is fulfilled if the LP-matrix is totally unimodular (or if each maximal quadratic submatrix belonging to a feasible basis has determinant  $\pm 1$ ). So, such a version of a UBTV would yield sharp upper bounds for the number of basic feasible solutions of such LP's.

It is well known [5], that for  $d \geq 3$  every cyclic  $d$ -polytope with  $v$  vertices has a vertex whose vertex figure (or *link*) is isomorphic to the boundary complex of a cyclic  $(d - 1)$ -polytope with  $v - 1$  vertices. It follows from this fact and the results of [8] that a smooth toric  $d$ -variety arising from a complete regular fan which is isomorphic to the boundary-complex of a cyclic  $d$ -polytope with  $v$  vertices possesses a smooth toric  $(d - 1)$ -subvariety which comes from a complete regular fan which is isomorphic to the boundary-complex of a cyclic  $(d - 1)$ -polytope with  $v - 1$  vertices. This fact and our Theorem directly imply:

COROLLARY 2. *Let  $\Delta$  be a complex which is isomorphic to the boundary-complex of the cyclic  $d$ -polytope with  $n$  vertices,  $n \geq d + 3 \geq 7$ . Then there does not exist a complex  $\Delta'$  with the properties (1) - (5) which is combinatorially equivalent to  $\Delta$ .*

This proves that for  $n \geq d + 3 \geq 7$ , no complete smooth toric  $d$ -variety has an underlying cone-system stemming from a cyclic  $d$ -polytope with  $n$  vertices.

Applying the methods of Section 3 to the three non-cyclic neighborly 3-spheres with 8 vertices given in the list of Grünbaum & Sreedharan [12], we were able to extend Corollary 1 from seven to eight vertices. Note that among these three spheres only two are boundary complexes of 4-polytopes with 8 vertices ( $P_{35}^8$  and  $P_{36}^8$  in [12]) while the third one is the non-polytopal *Brückner sphere* (denoted  $\mathcal{M}$  in [12]). More precisely, we have:

REMARK 3. *Let  $X$  be a complete toric 4-variety with  $\text{rk } H^2(X, \mathbb{Q}) = 4$ . Then  $\text{rk } H^4(X, \mathbb{Q}) \leq 10$ , and this bound is attained by four combinatorial types of fans. If, in addition,  $X$  is assumed to be smooth, then we have  $\text{rk } H^4(X, \mathbb{Q}) \leq 9$ .*

In view of these results we conjecture that for  $d \geq 4$ , no neighborly  $d$ -polytope with  $d + 3$  or more vertices gives rise to a complete smooth toric  $d$ -variety.

#### 4. References.

- [1] V.V. BATYREV, *Toroidal Fano 3-folds*, Math. USSR Izvestija, 19, No. 1 (1982), pp. 13–25.
- [2] V.I. DANILOV, *The geometry of toric varieties*, Math. USSR Izvestija, 17 (1981), pp. 97–154.
- [3] G. EWALD, *Spherical complexes and nonprojective toric varieties*, Discrete Comput. Geometry, 1 (1986), pp. 115–122.
- [4] G. EWALD, *Unsolved problems in combinatorial convexity relevant for algebraic geometry*, Abstracts of the 4th Geometry Symposium in Siegen (1986).
- [5] B. GRÜNBAUM, *Convex polytopes*, Interscience Publishers, New York, 1967.
- [6] V.L. KLEE AND P. KLEINSCHMIDT, *Geometry of the Gass-Saaty parametric LP algorithm*, Discrete Comput. Geometry (to appear).
- [7] P. KLEINSCHMIDT, *A classification of toric varieties with few generators*, Aequationes Mathematicae (to appear).
- [8] T. ODA, *Torus embeddings and applications*, Tata Inst. Fund Res. Lecture on Math. and Phys., 57 (Springer 1978).
- [9] R.P. STANLEY, *The upper bound conjecture and Cohen-Macaulay-rings*, Studies in Applied Math., 54 (1975), pp. 135–142.
- [10] R.P. STANLEY, *The number of faces of a simplicial convex polytope*, Advances in Math., 45 (1980), pp. 236–238.
- [11] B. STURMFELS, *Computational Synthetic Geometry*, Ph.D. Dissertation, University of Washington, 1987.
- [12] B. GRÜNBAUM AND V.P. SREEDHARAN, *An enumeration of simplicial 4-polytopes with 8 vertices*, J. Combinatorial Theory, 2 (1967), pp. 437–465.