

**ON SOME NEW EXACT SOLUTIONS OF NONLINEAR  
D'ALLEMBERT AND HAMILTON EQUATIONS**

By

**Wilhelm I. Fushchich**

and

**Renat Zhdanov**

**IMA Preprint Series # 468**

December 1988

# ON SOME NEW EXACT SOLUTIONS OF NONLINEAR D'ALLEMBERT AND HAMILTON EQUATIONS

BY WILHELM I. FUSHCHICH AND RENAT ZHDANOV†

**Abstract.** Some new exact solutions of d'Allembert-Hamilton and d'Allembert equations are obtained. The necessary conditions of integrability of over-determined d'Allembert-Hamilton system of nonlinear differential equations are established.

1. It was the Euler's idea (1734-1740 y.) that problem of integrating partial differential equations (PDE) could be solved by reducing them to ordinary equations (ODE). But one can not apply this idea to arbitrary PDE. Therefore it was suggested by Fushchich (1981, 1983) to restrict oneself by PDE possessing wide symmetry groups. This program was realized for some nonlinear wave equations by Fushchich and Serov (1983), Fushchich and Shtelen (1982) and Fushchich and Zhdanov (1987) (see also Beckers et al (1977), Patera et al (1975) and Grundland et al (1984). The vast list of references on this point can be found in Fushchich and Nikitin (1986).

When reducing PDE to ODE one has always to deal with the problem of investigating compatibility of some systems of PDE. For example, nonlinear d'Allembert equation

$$(1) \quad \square u = F_1(u), \quad \square = \partial_{x_0}^2 - \partial_{x_1}^2 - \partial_{x_2}^2 - \partial_{x_3}^2$$

with the aid of ansatz (Fushchich (1981))

$$(2) \quad u = \varphi(\omega), \quad \omega = \omega(x_0, x_1, x_2, x_3),$$

is reduced to the ODE having variable coefficients (Fushchich and Serov (1983))

$$(3) \quad \omega_\mu \omega^\mu \ddot{\varphi} + \square \omega \dot{\varphi} = F_1(\varphi),$$

where  $\omega_\mu \equiv \frac{\partial \omega}{\partial x^\mu}$ ,  $\mu = \overline{0, 3}$ ,  $\dot{\varphi} \equiv \frac{d\varphi}{d\omega}$ . Hereafter the summation over repeated indices in Minkowsky space having the metric  $g_{\mu\nu} = \text{diag}(1, -1, -1, -1)$  is supposed, i.e.  $\omega_\mu \omega^\mu \equiv g^{\mu\nu} \omega_\mu \omega_\nu = \omega_0^2 - \omega_1^2 - \omega_2^2 - \omega_3^2$ .

We demand new variable  $\omega$  to satisfy d'Allembert and Hamilton equations simultaneously

$$(4) \quad \square \omega = F_2(\omega),$$

$$(5) \quad \omega_\mu \omega^\mu = F_3(\omega).$$

---

†Institute of Mathematics, Repin Street 3, Kiev-4, USSR

As a result equation (3) takes the form

$$(6) \quad F_3(\omega\ddot{\varphi} + F_2(\omega)\dot{\varphi} = F_1(\varphi).$$

Winternitz and collaborators (see Beckers et al (1977), Patera et al (1975) and Patera et al (1976)) construct new variables  $\omega$  by using subgroup structure of the Poincaré group  $P(I, 3)$ . One can be easily convinced that invariants obtained in this way satisfy system (4), (5).

So to obtain set of variables  $\omega$  making possible to reduce multi-dimensional PDE (I) to ODE one has to consider the problem of compatibility of system (4), (5) and then to integrate it.

In the present paper compatibility of equations (4), (5) is investigated, i.e. all smooth functions ensuring the compatibility of d'Allembert-Hamilton system are described.

The direct application of Cartan's method of investigation of compatibility of over-determined PDE (Cartan (1946)) is rather difficult. To avoid arising difficulties we essentially use symmetry properties of system (4), (5) (Fushchich and Serov (1983) and Fushchich and Shtelen (1982)).

System (4), (5) via the change of dependent variable  $z = z(\omega)$  can be reduced to the following system

$$(7) \quad \square\omega = F(\omega),$$

$$(8) \quad \omega_\mu\omega^\mu = \lambda, \quad \lambda = \text{const.},$$

ODE (6) taking the form

$$(9) \quad \lambda\ddot{\varphi} + F(\omega)\dot{\varphi} = F_1(\varphi).$$

Before formulating the principal result of the paper we adduce without proof some auxiliary statements.

LEMMA I. *Solutions of system (7), (8) satisfy the identities*

$$(10) \quad \begin{aligned} \omega_{\mu\nu_1}\omega^{\mu\nu_1} &= -\lambda\dot{F}(\omega), \\ \omega_{\mu\nu_1}\omega^{\nu_1\nu_2}\omega_{\nu_2}^\mu &= \frac{1}{2!}\lambda^2\dot{F}(\omega), \\ \omega_{\mu\nu_1}\omega^{\nu_1\nu_2}\ddot{\omega}^{\nu_n\mu} &= \frac{1}{n!}(-\lambda)^n \frac{d^n F(\omega)}{d\omega^n}, \quad n \geq 0, \end{aligned}$$

where  $\omega_{\alpha\beta} \equiv \frac{\partial^2 \omega}{\partial x_\alpha \partial x_\beta}$ ,  $\alpha, \beta = \overline{0, 3}$ .

LEMMA 2. *Solutions of the system (7), (8) satisfy the following equality:*

$$(10') \quad \det(\omega_{\mu\nu}) = 0.$$

Let us now formulate the principal statement.

THEOREM 1. The necessary condition of compatibility of overdetermined system (7), (8) is as follows

$$(11) \quad F(\omega) = \begin{cases} 0, \\ \lambda(\omega + C_1)^{-1} \\ 2\lambda(\omega + C_1)[(\omega + C_1)^2 + C_2]^{-1}, \\ 3\lambda((\omega + C_1)^2 + C_2)[(\omega + C_1)^3 + 3C_2(\omega + C_1) + C_3]^{-1}, \end{cases}$$

where  $C_1, C_2, C_3$ -arbitrary constants.

Proof. By direct (and rather tiresome) verification one can be convinced that the following identity holds

$$(12) \quad \begin{aligned} &6(\omega_{\mu\nu_1}\omega^{\nu_1\nu_2}\omega_{\nu_2\nu_3}\omega^{\nu_3\mu}) - 8(\omega_\mu^\mu)(\omega_{\mu\nu_1}\omega^{\nu_1\nu_2}\omega_{\nu_2}^\mu) - \\ &- 3(\omega_{\mu\nu_1}\omega^{\mu\nu_1})^2 + 6(\omega_\mu^\mu)^2(\omega_{\mu\nu_1}\omega^{\mu\nu_1}) - (\omega_\mu\omega^\mu)^4 = 24\det(\omega_{\mu\nu}). \end{aligned}$$

Substituting (10), (10') into (12) one obtains nonlinear ODE for  $F(\omega)$

$$(13) \quad \lambda^3 \ddot{F} + 4\lambda^2 F \ddot{F} + 3\lambda^2 \dot{F}^2 + 6\lambda \dot{F} F^2 + F^4 = 0,$$

where  $\dot{F} \equiv \frac{dF}{d\omega}$ .

General solution of equation (13) is given by formulae (11). Theorem is proved.

Note 1. Compatibility of three-dimensional d'Allembert-Hamilton system has been investigated in detail by Collins (1976). Collins essentially used geometrical methods which could not be generalized to higher dimensions.

Using Lie's method (see e.g. Olver (1986)) one can prove the following statement.

THEOREM 2. The sytem of PDE (7), (8) is invariant under the 15 - parameter conformal group  $C(1, 3)$  iff

$$(14) \quad F(\omega) = 3\lambda(\omega + C)^{-1}, \quad \lambda > 0, \quad C = const.$$

Note 2. Formula (14) can be obtained from (11) by putting  $C_2 = C_3 = 0$ . So Theorem 2 demonstrates close connection between compatibility of a system of PDE and its symmetry.

Note 3. It is common knowledge that PDE (7) is invariant under the group  $C(I, 3)$  iff  $F(\omega) = \lambda\omega^3$  (Fushchich and Serov (1983)). Consequently, an additional constraint (8) changes essentially symmetry properties of d'Allenbert equation (choosing  $F_3(\omega)$  in a proper way one can obtain conformally-invariant system of the form (4), (5) under arbitrary  $F_2(\omega)$ ).

2. Let us list explicit form of some exact solutions of d'Allembert-Hamilton system and reduced ODE for function  $\varphi(\omega)$ .

N <sup>o</sup>	$\lambda$	$F(\omega)$	$\omega = \omega(x)$	ODE for $\varphi(\omega)$
1.	1	0	$a_\mu x^\mu$	$\ddot{\varphi} = F_1(\varphi)$
2.	1	$\omega^{-1}$	$[(a_\mu x^\mu)^2 - (b_\mu x^\mu)^2]^{1/2}$	$\dot{\varphi} + \omega^{-1}\dot{\varphi} = F_1(\varphi)$
3.	1	$2\omega^{-1}$	$[(a_\mu x^\mu)^2 - (b_\mu x^\mu)^2 - (c_\mu x^\mu)^2]^{1/2}$	$\ddot{\varphi} + 2\omega^{-1}\dot{\varphi} = F_1(\varphi)$
4.	1	$3\omega^{-1}$	$(x_\mu x^\mu)^{1/2}$	$\ddot{\varphi} + 3\omega^{-1}\dot{\varphi} = F_1(\varphi)$
5.	-1	0	$(b_\mu x^\mu) \cos(h_1) + (c_\mu x^\mu) \sin(h_1) + g_1$ $a_\mu x^\mu - (b_\mu x^\mu) \cos(h_2) -$ $-(c_\mu x^\mu) \sin(h_2) - g_2 = 0$	$\ddot{\varphi} = -F_1(\varphi)$
6.	-1	$-\omega^{-1}$	$[(b_\mu x^\mu + h_1)^2 + (c_\mu x^\mu + h_2)^2]^{1/2}$	$\ddot{\varphi} + \omega^{-1}\dot{\varphi} = -F_1(\varphi)$
7.	-1	$-2\omega^{-1}$	$[(b_\mu x^\mu)^2 + (c_\mu x^\mu)^2 + (d_\mu x^\mu)^2]^{1/2}$	$\ddot{\varphi} + 2\omega^{-1}\dot{\varphi} = -F_1(\varphi)$
8.	0	0	$h_1$	$0 = F_1(\varphi)$

Here  $h_1, g_1$ -arbitrary smooth functions on  $a_\mu x^\mu + d_\mu x^\mu$ ,  $h_2, g_2$ -on  $\omega + d_\mu x^\mu$ ;  $a_\mu, b_\mu, c_\mu, d_\mu$ -arbitrary real parameters satisfying conditions of the form

$$-a_\mu a^\mu = b_\mu b^\mu = c_\mu c^\mu = d_\mu d^\mu = -1,$$

$$a_\mu b^\mu = a_\mu c^\mu = a_\mu d^\mu = b_\mu c^\mu = b_\mu d^\mu = c_\mu d^\mu = 0.$$

3. Natural generalization of the formula (2) is given by ansatz of the form (Fushchich (1981))

$$(15) \quad u(x) = f(x)\varphi(\omega).$$

Some multi-parameter families of exact solutions of nonlinear d'Allembert equation with nonlinearity  $F_1 = \tau u^k$ ,  $\tau, k$ -const., were constructed with the help of ansatz (15) by Fushchich and Serov (1983).

Omitting intermediate calculations we write down new family of solutions of equation (I) under  $F_1 = \tau u^k$  obtained via ansatz (15)

$$u(x) = R^{-1} [C_6 + \frac{1}{2}\tau(1-k)^2 \int R^{1-k}(\omega)d\omega]^{1/(1-k)} \times$$

$$\times \left\{ \frac{1}{2}(a_\mu x^\mu - d_\mu x^\mu) - \frac{1}{2}\dot{R}R^{-1}[(b_\mu x^\mu)^2 + (c_\mu x^\mu)^2] + \right.$$

$$\left. + f_0[(b_\mu x^\mu)^2 - (c_\mu x^\mu)^2] + f_1 b_\mu x^\mu + f_2 \right\}^{1/(1-k)},$$

where

$$\begin{aligned}
 f_0(\omega) &= \frac{1}{2}C_1R^{-2}(\omega), \\
 f_1(\omega) &= \frac{1}{2}C_4R(\omega)\exp\{4C_1\int R^{-2}(\omega)d\omega\}, \\
 f_2(\omega) &= C_4\int R(\omega)\exp\{4C_1\int R^{-2}(\omega)d\omega\}d\omega + C_5 \\
 \text{and } \omega &= a_\mu x^\mu + d_\mu x^\mu, \quad C_1, \dots, C_6 - \text{const.}
 \end{aligned}$$

Function  $R = R(\omega)$  is determined by formulae

$$R(\omega) = \begin{cases} \frac{\varepsilon}{2}[(C_2\omega + C_3)^2 - 16C_1^2]^{\frac{1}{2}}, \\ (8\varepsilon C_1\omega + C_2)^{\frac{1}{2}}, \quad \varepsilon = \pm 1. \end{cases}$$

#### REFERENCES

- [1] BECKERS J., PATERA J. AND WINTERNITZ P., *J. Math. Phys.*, 18 (1977), pp. 72-83.
- [2] CARTAN E., *Les systèmes différentiels extérieurs et leur applications scientifiques*, Paris: Hermann, 1946.
- [3] COLLINS, C.B., *Proc. Cambr. Phil. Soc.*, 80 (1976), pp. 165-172.
- [4] FUSHCHICH, W.I., *The symmetry of mathematical physics problems, in Algebraic-theoretical studies in mathematical physics*, Kiev Institute of Mathematics (1981), pp. 6-28.
- [5] —————, *On symmetry and some exact solutions of some many-dimensional equations of mathematical physics, in Theoretical-algebraic methods in mathematical physics problems*, Kiev Institute of Mathematics (1983), pp. 4-23.
- [6] FUSHCHICH, W.I. AND NIKITIN, A.G., *Symmetries of Maxwell's Equations*, Dordrecht: D. Reidel Publ. Comp., 1987.
- [7] FUSHCHICH, W.I. AND SEROV, N.I., *The symmetry and some exact solutions of the nonlinear multidimensional Liouville, d'Alambert and eikonal equations*, *J. Phys. A: Math. Gen.*, 16 (1983), pp. 3645-3656.
- [8] FUSHCHICH, W.I. AND SHTELEN, W.M., *Conformal symmetry and new exact solutions of  $SU_2$  Yang-Mills theory*, *Lett. Nuovo Cim.*, 34 (1982), pp. 498-502.
- [9] FUSHCHICH, W.I. AND ZHDANOV, R.Z., *On some exact solutions of a system of non-linear differential equations for spinor and vector fields*, *J. Phys. A: Math. Gen.*, 20 (1987), pp. 4173-4190.
- [10] OLVER, P.J., *Applications of Lie Groups to differential equations*, Springer-Verlag, New York, 1986.
- [11] PATERA, J., WINTERNITZ, P. AND ZASSENHAUS, H., *Continuous subgroups of fundamental group of physics I. General Method and the Poincare group*, *J. Math. Phys.*, 16 (1975), pp. 1597-1624.
- [12] GRUNDLAND, A.M., HARMAD, J. AND WINTERNITZ, P., *Symmetry reduction for nonlinear relativistically invariant equations*, *Jour. Math. Phys.*, 25 (1984), pp. 491-806.